





- A. Breast
- **B.** <sup>1</sup>/<sub>2</sub> oz. Liquid Lubricant
- C. 20 ml Syringe with Tubing

1. Phaladalad

#### **General Instructions for Use**

- 1. Ensure the stopcock is connected to syringe and small tubing.
- 2. Fill the syringe with water.
- **3.** Dip the tubing into the  $\frac{1}{2}$  oz. lubricant provided. (See figure 1.)
- Place the lubricated tubing into nipple hole.
- 5. Slowly depress the syringe until you see the breast swell slightly. (See figure 2.)
- Please see reverse side for care.



Life form.

Year





The maximum capacity of fluid in breast is 12 cc.







## Mastitis LF00861U

An inflamed breast due to bacterial infection on a damaged nipple.

### Flat LF008621

Nipple is not protruding outward even when stimulated

# **Inverted LF00860U**

The nipple is protruding inward instead of outward. The nipple is inward even when stimulated.

# **Care of Breast Trainers**

- Wash breasts with warm soap and water.
- 2. Make sure breasts are dry before storing.
- 3. Baby powder can be used to add softness.
- 4. Excessive rubbing of breasts can damage the trainer.
- 5. Handle the breast replica like you would real breasts.
- Contact with ink, printed materials, and ballpoint pen will stain breast replica.
- 7. Ensure the lactation breast is empty of fluid before storing.

### Set of 4 Breast Conditions LF00864I

